

Please click price below and you will be redirected to our pharmacy.

## Isordil Sublingual



Generic **Isordil Sublingual** is used for preventing symptoms of angina (chest pain) caused by heart disease. This medicine is used alone or with other medicines. This medicine is NOT intended for the immediate relief of acute attacks of angina.

Active ingredient: [Isosorbide Dinitrate](#)

**Brand(s):**Sorbitrate

**Manufacturer:** Piramal Enterprise Ltd.

**Disease(s):**Angina

### Isordil Sublingual 5mg

	Per pill	Per Pack	Order
50 tablet	\$ 0.90	<b>\$ 45.00</b>	<a href="#">Add to cart</a>
100 tablet	\$ 0.85	<b>\$ 85.00</b> <del>\$90.00</del>	<a href="#">Add to cart</a>
150 tablet	\$ 0.80	<b>\$ 120.00</b> <del>\$135.00</del>	<a href="#">Add to cart</a>
200 tablet	\$ 0.75	<b>\$ 150.00</b> <del>\$180.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
250 tablet	\$ 0.70	<b>\$ 175.00</b> <del>\$225.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
300 tablet	\$ 0.65	<b>\$ 195.00</b> <del>\$270.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Isosorbide Dinitrate Sublingual tablet

### What is this medicine?

ISOSORBIDE DINITRATE is a type of vasodilator. It relaxes blood vessels, increasing the blood and oxygen supply to your heart. This medicine is used to prevent chest pain caused by angina. It will not help to stop an episode of chest pain.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- previous heart attack or heart failure
- an unusual or allergic reaction to isosorbide dinitrate, nitrates, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I use this medicine?

Place this medicine under the tongue. Do not cut, crush or chew this medicine. Take this medicine 15 minutes before an activity that may cause chest pain or as directed. Do not take your medicine more often than directed.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.

