

Please click price below and you will be redirected to our pharmacy.

## Clopidogrel



**Clopidogrel** is used for reducing the risk of stroke or heart attack in patients who have already had a heart attack or stroke or have other circulatory problems due to narrowing and hardening of the arteries.

Active ingredient: [Clopidogrel](#)

**Brand(s):**Caplor / Clavix

**Manufacturer:** Ind-Swift Laboratories Ltd. / Intas Pharmaceuticals Ltd.

**Disease(s):**Stroke / Heart Attack

### Clopidogrel 75mg

	Per pill	Per Pack	Order
30 tablet	\$ 0.96	<b>\$ 28.95</b>	<a href="#">Add to cart</a>
60 tablet	\$ 0.85	<b>\$ 50.74</b> <del>\$57.90</del>	<a href="#">Add to cart</a>
90 tablet	\$ 0.82	<b>\$ 74.04</b> <del>\$86.85</del>	<a href="#">Add to cart</a>
120 tablet	\$ 0.79	<b>\$ 94.94</b> <del>\$115.80</del>	<a href="#">Add to cart</a>
180 tablet	\$ 0.76	<b>\$ 136.75</b> <del>\$173.70</del>	<a href="#">Add to cart</a>
360 tablet	\$ 0.73	<b>\$ 262.15</b> <del>\$347.40</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Clopidogrel tablet

### What is this medicine?

**CLOPIDOGREL** helps to prevent blood clots. It reduces the chance of having a heart attack or a stroke in people who have already had a heart attack or a stroke. **Clopidogrel** can also decrease the chance of a heart attack or stroke in certain groups of people at high risk for these events.

### What should my health care professional know before I take this medicine?

They need to know if you have any of the following conditions:

- bleeding disorder or hemophilia
- liver disease
- recent surgery or trauma
- stomach or intestinal ulcers
- an unusual or allergic reaction to **clopidogrel**, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I take this medicine?

Take **clopidogrel** tablets by mouth. Swallow the tablets with a drink of water. Do not take your medicine more often than directed.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.

