

Please click price below and you will be redirected to our pharmacy.

## Eulexin



**Generic Eulexin** is an antiandrogen. It works in the body to prevent the actions of androgens (male hormones). It is used in the treatment of prostate cancer.

Active ingredient: [Flutamide](#)

**Brand(s):**Cytomid / Flutide

**Manufacturer:** Cipla Limited / Samarth Pharma Pvt.Ltd.

**Disease(s):**Prostate Cancer

### Eulexin 250mg

	Per pill	Per Pack	Order
30 tablet	\$ 2.67	<b>\$ 79.95</b>	<a href="#">Add to cart</a>
60 tablet	\$ 2.50	<b>\$ 149.95</b> <del>\$159.90</del>	<a href="#">Add to cart</a>
90 tablet	\$ 2.22	<b>\$ 199.95</b> <del>\$239.85</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 tablet	\$ 2.08	<b>\$ 249.95</b> <del>\$319.80</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
180 tablet	\$ 1.83	<b>\$ 329.95</b> <del>\$479.70</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
360 tablet	\$ 1.61	<b>\$ 579.95</b> <del>\$959.40</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

### Flutamide tablet

#### What is this medicine?

FLUTAMIDE blocks the effect of the male hormone called testosterone. It is used in combination with another medication to treat prostate cancer.

#### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- cigarette smoker
- glucose-6-phosphate dehydrogenase deficiency
- hemoglobin M disease
- lactose intolerance
- liver disease
- an unusual or allergic reaction to flutamide, other chemotherapy, other medicines, foods, dyes, or preservatives
- your partner is pregnant or trying to get pregnant

#### How should I use this medicine?

Take this medicine by mouth with a glass of water. Do not take your medicine more often than directed. Finish the full course prescribed by your doctor or health care professional, even if the capsules make you feel sick. Do not stop taking except on the advice of your doctor or health care professional.

Talk to your pediatrician regarding the use of this medicine in children.

