

Please click price below and you will be redirected to our pharmacy.

## Addyi



Addyi is used to treat decreased sexual desire in women who have not gone through menopause and who have never had low sexual desire in the past.

Active ingredient: [Flibanserin](#)

Brand(s): Fliban

Manufacturer: Centurion Laboratories Private Limited

Disease(s): Decreased Sexual Desire / Low Libido

### Addyi 100mg

	Per pill	Per Pack	Order
12 tablet	\$ 8.33	<b>\$ 99.95</b>	<a href="#">Add to cart</a>
20 tablet	\$ 8.00	<b>\$ 159.95</b> <del>\$166.58</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
32 tablet	\$ 6.25	<b>\$ 199.95</b> <del>\$266.53</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
60 tablet	\$ 5.83	<b>\$ 349.95</b> <del>\$499.75</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
92 tablet	\$ 5.65	<b>\$ 519.95</b> <del>\$766.28</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 tablet	\$ 5.42	<b>\$ 649.95</b> <del>\$999.50</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Flibanserin tablet

### What is this medicine?

FLIBANSERIN is used to treat decreased sexual desire in women who have not gone through menopause and who have never had low sexual desire in the past. Flibanserin is for use only when low sexual desire is NOT caused by a medical condition, a mental disorder, relationship problems, or by using drugs or other medicines.

Flibanserin is not for women who have already gone through menopause. This medicine is also not for use by men.

### What should I tell my health care provider before I take this medicine?

**You must not drink alcohol while taking flibanserin. It may cause you to have dangerously low blood pressure.**

You should not use flibanserin if you are allergic to it, or if you have:

- low blood pressure;
- a history of alcoholism or drug addiction;
- liver disease;
- a history of depression or mental illness; or
- if you currently drink alcohol.

Some medicines can cause unwanted or dangerous effects when used with flibanserin. Your doctor may need to change your treatment plan if you use any of the following drugs:

